

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

SRVs in the Solar Neighbourhood

I.S. Glass

*South African Astronomical Observatory, PO Box 9, Observatory 7935,
South Africa*

Abstract.

Period-luminosity sequences have been shown to exist among the Semi-Regular Variables (SRVs) of the Magellanic Clouds (Wood et al, 1999), the Bulge of the Milky Way galaxy (Glass & Schultheis, 2003) and elsewhere. It would clearly be useful to have absolute (trigonometric) calibrations of these relations. This paper investigates whether the sequences can be seen among the M-type giant SRVs of the solar neighbourhood. Mass loss phenomena among these stars and their dependence on period and spectral sub-type are also discussed.

The periods of 57 bright M-type SRVs have been extracted from Percy, Wilson & Henry (2001), Percy, Dunlop & Kassim (2001), Percy, Nyssa & Henry (2001) and Percy et al (2004). These have been supplemented with information from Bedding & Zijlstra (1998) and CDS, Strasbourg. Parallaxes have been taken from the Hipparcos Catalogue, via CDS, and K magnitudes from Neugebauer & Leighton (1969). Fig 1 shows the resulting M_K , $\log P$ diagram. Many stars have more than one period and all have been plotted. The approximate positions of the sequences seen in the Bulge are given for comparison.

The scatter in Fig 1 is rather large, but one can almost convince oneself that the A and B sequences are separately visible. The D certainly is. The C sequence, occupied by Miras, is not expected to be populated because of the choice of sample; nevertheless a few objects do appear to lie on it. It is hoped that more extensive photometric observations will in due course lead to better magnitudes and periods and that improved parallax observations will become available, so that the sequences will eventually be more precisely defined.

IRAS 12 μ m magnitudes have been combined with those at K to form $K - [12]$ colours and these have been plotted against M sub-type in Fig 2 and against $\log P$ in Fig 3. As in the case of the NGC6522 field, infrared excesses (due to mass-loss) appear in M-K types later than about M4 and at periods longer than about 70 days (cf Alard et al, 2001, Glass & Schultheis, 2002).

I thank Dr T. Lloyd Evans, Dr M. Schultheis and Mr N. Matsunaga for useful comments during this work.

References

- Alard, C. et al, 2001, ApJ, 552, 2001
- Bedding, T.R., Zijlstra, A.A. 1998, ApJ, 506, L47
- Glass, I.S., Schultheis, M. 2003, MNRAS, 345, 39
- Glass, I.S., Schultheis, M. 2002, MNRAS, 337, 519
- Neugebauer, G., Leighton, R.B. 1969, Two-Micron Sky Survey, NASA SP-3047, Washington, D.C.
- Percy, J.R., Bakos, A.G., Besla, G., Hou, D., Velocci, V., Henry, G.W. 2004, IAU Coll. 193, ASP Conf. Ser., 310, p348




Figure 1. M_K , $\log P$ diagram for SRVs in the solar neighbourhood. The positions of the Galactic Bulge A,B,C and D sequences from Glass & Schultheis (2003) are shown. Stars plotted with light symbols have $K - [12] > 1.0$, indicative of dust shells. Some interesting outliers have been labelled.




Figure 2. *Left:* $K - [12]$ colour vs M-K type for the local SRVs. Note the prevalence of significant excess $12\ \mu\text{m}$ radiation beyond types M3-M4. *Right:* $K - [12]$ colour vs log period for local SRVs. All periods are shown. The long-period D sequence stars usually do not exhibit $12\ \mu\text{m}$ excess.

- Percy, J.R., Nyssa, Z., Henry, G.W. 2001, IBVS 5209
Percy, J.R., Wilson, J.B. and Henry, G.W. 2001, PASP, 113, 983
Percy, J.R., Dunlop, H., Kassim, L., Thompson, R.R. 2001, IBVS 5041
Wood, P.R. and the MACHO Team 1999, in IAU Symp. 191, p151